

QUANTUM TOROIDAL ALGEBRAS AND THEIR VERTEX REPRESENTATIONS

YOSHIHISA SAITO

ABSTRACT. We construct the vertex representations of the quantum toroidal algebras $U_q(\mathfrak{sl}_{n+1,tor})$. In the classical case the vertex representations are not irreducible. However in the quantum case they are irreducible.

For $n=1$, we construct a set of finitely many generators of $U_q(\mathfrak{sl}_{2,tor})$.

1. INTRODUCTION

The classical toroidal algebras have been studied by many authors [MEY], [S], [Y], etc. Here “classical” means $q = 1$. The definition of the quantum toroidal algebras is given in [GKV]. They gave a geometric realization of the quantum toroidal algebras without any results on their representation theory. Recently Varagnolo and Vasserot [VV] proved a Schur-type duality between representations of the quantum toroidal algebras and the double affine Hecke algebra introduced by Cherednik [C]. This is an analogue of the duality between the quantum affine algebras and the affine Hecke algebras given by Chari and Pressly [CP]. In [VV] only the representations of “trivial central charge” was studied. It is known that there are two subalgebras $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ and $U_q^{(2)}(\widehat{\mathfrak{sl}}_{n+1})$ of $U_q(\mathfrak{sl}_{n+1,tor})$ which are isomorphic to $U_q(\widehat{\mathfrak{sl}}_{n+1})$. We say the $U_q(\mathfrak{sl}_{n+1,tor})$ -module M has the trivial central charge if M has level 0 as both $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module and $U_q^{(2)}(\widehat{\mathfrak{sl}}_{n+1})$ -module. It is an analogue of level 0 representations of the affine quantum algebras. In this paper we say that M has a level $(0, 0)$ instead of the trivial central charge. The first 0 means that M has a level 0 as a $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module and the second 0 means that M has a level 0 as a $U_q^{(2)}(\widehat{\mathfrak{sl}}_{n+1})$ -module.

In this paper we try to consider an analogue of “integrable representations” in the toroidal case. Let us recall the integrability of quantum Kac-Moody modules. Let $U_q(\mathfrak{g})$ be a quantum Kac-Moody algebra and V a $U_q(\mathfrak{g})$ -module. We say V is integrable if V has a weight space decomposition and locally nilpotent actions of the Chevalley generators of

The author is supported by the JSPS Research Fellowships for Young Scientists.

$U_q(\mathfrak{g})$. Therefore the definition of “integrability” needs Chevalley-type generators. The toroidal algebras is defined through Drinfeld type of generators and its Chevalley-type generators are not known. Therefore we are not able to define the integrability at this moment. However, in the affine case, Frenkel-Jing [FJ] realized the integrable representations with level 1 by the vertex representations. Thus if there are “vertex representations” of quantum toroidal algebras, they must be interesting examples of the integrable representations still not defined. In the $q = 1$ case, vertex representations of the toroidal algebras have been already considered by Moody-Eswara Rao-Yokonuma [MEY]. In this paper for we construct the q -analogue of the representations defined by them for $\mathfrak{g} = \mathfrak{sl}_{n+1}$ with level $(1, 0)$ and $(1, 1)$. Therefore we give a new class of the representations of the quantum toroidal algebras. In the $q = 1$ case, the Fock modules are not irreducible over the Heisenberg algebra and the vertex representations are not irreducible over the toroidal algebra. In the quantum case, it is not the case: the Fock modules are irreducible and also the vertex representations with level $(1, 0)$ are irreducible.

The algebra $U_q(\mathfrak{g}_{tor})$ has infinitely many generators satisfying infinitely many relations (See §2). It is preferable that $U_q(\mathfrak{g}_{tor})$ is written by finitely many generators with finitely many relations. According to [GKV] there are finitely many generators of $U_q(\mathfrak{sl}_{2,tor})$ but the relations among these generators are highly non-trivial. In this paper we give an explicit form of finitely many generators of $U_q(\mathfrak{sl}_{2,tor})$ and closed relations of them (See §4). They coincide with the generators by Vasserot [V].

2. DEFINITION OF QUANTUM TOROIDAL ALGEBRAS

2.1. Notations. Let \mathfrak{g} be a complex semisimple Lie algebra of type A_n and $\hat{\mathfrak{g}}$ an affine Kac-Moody Lie algebra of type $A_n^{(1)}$. We denote their Cartan subalgebras by \mathfrak{h} and $\hat{\mathfrak{h}}$ respectively. We denote by $\bar{\alpha}_1, \dots, \bar{\alpha}_n$ the simple roots of \mathfrak{g} , by $\bar{h}_1, \dots, \bar{h}_n$ the simple coroots of \mathfrak{g} , by $\bar{\Lambda}_1, \dots, \bar{\Lambda}_n$ the fundamental weights of \mathfrak{g} , by $\alpha_0, \dots, \alpha_n$ the simple roots of $\hat{\mathfrak{g}}$, by h_0, \dots, h_n the simple coroots of $\hat{\mathfrak{g}}$ and $\Lambda_0, \dots, \Lambda_n$ the fundamental weights of $\hat{\mathfrak{g}}$. Let $\bar{Q} = \bigoplus_{i=1}^n \mathbb{Z}\bar{\alpha}_i$ be the root lattice of \mathfrak{g} , $\bar{P} = \bigoplus_{i=1}^n \mathbb{Z}\bar{\Lambda}_i$ the weight lattice of \mathfrak{g} , $Q = \bigoplus_{i=0}^n \mathbb{Z}\alpha_i$ the root lattice of $\hat{\mathfrak{g}}$ and $P = \bigoplus_{i=0}^n \mathbb{Z}\Lambda_i \oplus \mathbb{Z}\delta$ the weight lattice of $\hat{\mathfrak{g}}$. Here δ is the null root.

We denote the pairing of \mathfrak{h} and \mathfrak{h}^* (*resp.* $\hat{\mathfrak{h}}$ and $\hat{\mathfrak{h}}^*$) by $\langle \cdot, \cdot \rangle$. The invariant bilinear form on P is given by $(\alpha_i | \alpha_j) = -\delta_{ij-1} + 2\delta_{ij} - \delta_{ij+1}$

and $(\delta|\delta) = 0$. The projection form P to \overline{P} is given by $\overline{\Lambda}_i = \Lambda_i - \Lambda_0$ and $\overline{\delta} = 0$.

2.2. We will give the definition of the quantum toroidal algebra $U_q(\mathfrak{g}_{tor})$.

Definition 2.2.1. Let $M = (m_{ij})_{0 \leq i, j \leq n}$ be a skew-symmetric $(n+1) \times (n+1)$ -matrix with integral coefficients and let κ be an element of $\mathbb{Q}(q)^\times$. $U_q(\mathfrak{g}_{tor})$ is an associated algebra over $\mathbb{Q}(q)$ with generators :

$$E_{i,k}, F_{i,k}, H_{i,l}, K_i^\pm, q^{\pm \frac{1}{2}c}, q^{\pm d_1}, q^{\pm d_2},$$

for $k \in \mathbb{Z}$, $l \in \mathbb{Z} \setminus \{0\}$ and $i = 0, 1, \dots, n$.

We introduce $K_{i,k}^\pm$ as the Fourier components of the following generating functions:

$$K_i^+(z) = \sum_{k \geq 0} K_{i,k}^+ z^{-k} = K_i^+ \exp((q - q^{-1}) \sum_{k \geq 1} H_{i,k} z^{-k}),$$

$$K_i^-(z) = \sum_{k \leq 0} K_{i,k}^- z^{-k} = K_i^- \exp(-(q - q^{-1}) \sum_{k \geq 1} H_{i,-k} z^k).$$

The defining relations of $U_q(\mathfrak{g}_{tor})$ are then written as follows:

$$q^{\pm \frac{1}{2}c} \text{ are central,} \quad (2.2.1)$$

$$K_i^+ K_i^- = K_i^- K_i^+ = 1, \quad (2.2.2)$$

$$[K_i^\pm, K_j^\pm] = 0, \quad (2.2.3)$$

$$[K_i^\pm, H_{j,l}] = 0, \quad (2.2.4)$$

$$[H_{i,k}, H_{j,l}] = \delta_{k+l,0} \frac{1}{k} [k \langle h_i, \alpha_j \rangle] \frac{q^{kc} - q^{-kc}}{q - q^{-1}} \kappa^{-km_{ij}}, \quad (2.2.5)$$

$$[q^{\pm d_i}, K_j^\pm] = 0, \quad (2.2.6)$$

$$q^{d_1} H_{j,l} q^{-d_1} = q^l H_{j,l}, \quad (2.2.7)$$

$$[q^{\pm d_2}, H_{j,l}] = 0, \quad (2.2.8)$$

$$q^{d_1} E_{j,k} q^{-d_1} = q^k E_{j,k}, \quad (2.2.9)$$

$$q^{d_1} F_{j,k} q^{-d_1} = q^k F_{j,k},$$

$$q^{d_2} E_{j,k} q^{-d_2} = q^{\delta_{j0}} E_{j,k}, \quad (2.2.10)$$

$$q^{d_2} F_{j,k} q^{-d_2} = q^{-\delta_{j0}} F_{j,k},$$

$$K_i^+ E_{j,k} K_i^- = q^{\langle h_i, \alpha_j \rangle} E_{j,k}, \quad (2.2.11)$$

$$K_i^+ F_{j,k} K_i^- = q^{-\langle h_i, \alpha_j \rangle} F_{j,k},$$

$$[H_{i,k}, E_{j,l}] = \frac{1}{k} [k \langle h_i, \alpha_j \rangle] q^{-\frac{1}{2}|k|c} \kappa^{-km_{ij}} E_{j,k+l}, \quad (2.2.12)$$

$$[H_{i,k}, F_{j,l}] = -\frac{1}{k} [k \langle h_i, \alpha_j \rangle] q^{\frac{1}{2}|k|c} \kappa^{-km_{ij}} F_{j,k+l},$$

$$\kappa^{m_{ij}} E_{i,k+1} E_{j,l} - q^{\langle h_i, \alpha_j \rangle} \kappa^{m_{ij}} E_{j,l} E_{i,k+1} = q^{\langle h_i, \alpha_j \rangle} E_{i,k} E_{j,l+1} - E_{j,l+1} E_{i,k}, \quad (2.2.13)$$

$$\kappa^{m_{ij}} F_{i,k+1} F_{j,l} - q^{-\langle h_i, \alpha_j \rangle} \kappa^{m_{ij}} F_{j,l} F_{i,k+1} = q^{-\langle h_i, \alpha_j \rangle} F_{i,k} F_{j,l+1} - F_{j,l+1} F_{i,k},$$

$$[E_{i,k}, F_{j,l}] = \delta_{i,j} \frac{1}{q - q^{-1}} \{ q^{\frac{1}{2}(k-l)c} K_{i,k+l}^+ - q^{\frac{1}{2}(l-k)c} K_{i,k+l}^- \}, \quad (2.2.14)$$

$$\sum_{\sigma \in \mathfrak{S}_m} \sum_{r=0}^m (-1)^r \begin{bmatrix} m \\ r \end{bmatrix} E_{i,k_{\sigma(1)}} \cdots E_{i,k_{\sigma(r)}} E_{j,l} E_{i,k_{\sigma(r+1)}} \cdots E_{i,k_{\sigma(m)}} = 0, \quad (2.2.15)$$

$$\sum_{\sigma \in \mathfrak{S}_m} \sum_{r=0}^m (-1)^r \begin{bmatrix} m \\ r \end{bmatrix} F_{i,k_{\sigma(1)}} \cdots F_{i,k_{\sigma(r)}} F_{j,l} F_{i,k_{\sigma(r+1)}} \cdots F_{i,k_{\sigma(m)}} = 0,$$

for $i \neq$

j ,

where $m = 1 - \langle h_i, \alpha_j \rangle$.

In these relations we denote $[k] = \frac{q^k - q^{-k}}{q - q^{-1}}$, $[n]! = \prod_{k=1}^n [k]$, $\begin{bmatrix} m \\ r \end{bmatrix} = \frac{[m]!}{[r]![m-r]!}$.

2.3. Let $U_q''(\mathfrak{g}_{tor})$ be the subalgebra of $U_q(\mathfrak{g}_{tor})$ generated by $E_{i,k}, F_{i,k}, K_i^\pm, H_{i,l}, q^{\pm \frac{1}{2}c}$. Let $U_q^{\phi'}(\mathfrak{g}_{tor})$ (resp $U_q^{\phi}(\mathfrak{g}_{tor})$) be the subalgebra generated by $U_q''(\mathfrak{g}_{tor})$ and $q^{\pm d_1}$ (resp $q^{\pm d_2}$). Let $U_q^{(1)'}(\widehat{\mathfrak{sl}}_{n+1})$ be the subalgebra generated by $E_{i,k}, F_{i,k}, K_i^\pm, H_{i,l}, q^{\pm \frac{1}{2}c}$ ($1 \leq i \leq n, k \in \mathbb{Z}, l \in \mathbb{Z} \setminus \{0\}$) and $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ the subalgebra generated by $U_q^{(1)'}(\widehat{\mathfrak{sl}}_{n+1})$ and $q^{\pm d_1}$. Let $U_q^{(2)'}(\widehat{\mathfrak{sl}}_{n+1})$ the subalgebra generated by $E_{i,0}, F_{i,0}, K_i^\pm$ ($0 \leq i \leq n$) and $U_q^{(2)}(\widehat{\mathfrak{sl}}_{n+1})$ the subalgebra generated by $U_q^{(2)'}(\widehat{\mathfrak{sl}}_{n+1})$ and $q^{\pm d_2}$. By the definition $U_q^{(2)}(\widehat{\mathfrak{sl}}_{n+1})$ is isomorphic to $U_q(\widehat{\mathfrak{sl}}_{n+1})$ and $U_q^{(2)'}(\widehat{\mathfrak{sl}}_{n+1})$ is

isomorphic to $U'_q(\widehat{\mathfrak{sl}}_{n+1})$.

The following are straight forward.

Lemma 2.3.1. *For $1 \leq i \leq n$ let $\overline{E_{i,k}} = E_{i,k} \kappa^{\sum_{j=1}^i km_{j-1j}}$, $\overline{F_{i,k}} = F_{i,k} \kappa^{\sum_{j=1}^i km_{j-1j}}$, $\overline{H_{i,l}} = H_{i,l} \kappa^{\sum_{j=1}^i lm_{j-1j}}$. Then the relations between $\overline{E_{i,k}}$, $\overline{F_{i,k}}$, $\overline{H_{i,l}}$ and K_i^\pm are precisely the relations of Drinfeld generators of $U_q(\widehat{\mathfrak{sl}}_{n+1})$. That is, $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ is isomorphic to $U_q(\widehat{\mathfrak{sl}}_{n+1})$ and $U_q^{(1)'}(\widehat{\mathfrak{sl}}_{n+1})$ is isomorphic to $U'_q(\widehat{\mathfrak{sl}}_{n+1})$.*

Lemma 2.3.2. *Let $K_\delta^\pm = \prod_{i=0}^n K_i^\pm$. Then K_δ^\pm are the central elements of $U_q(\mathfrak{g}_{tor})$.*

Note that $q^{\pm c}$ is the central elements of $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ and K_δ^\pm the central elements of $U_q^{(2)}(\widehat{\mathfrak{sl}}_{n+1})$.

3. VERTEX REPRESENTATIONS

3.1. Heisenberg algebras. In this section we shall give the vertex representations of $U_q(\mathfrak{g}_{tor})$. We assume $c = 1$.

Consider a $\mathbb{Q}(q)$ -algebra S_n generated by $H_{i,l}$ ($0 \leq i \leq n$, $l \in \mathbb{Z} \setminus \{0\}$) satisfying:

$$[H_{i,k}, H_{j,l}] = \delta_{k+l,0} \frac{1}{k} [k \langle h_i, \alpha_j \rangle] \frac{q^k - q^{-k}}{q - q^{-1}} \kappa^{-km_{ij}}. \quad (3.1.1)$$

We call S_n the Heisenberg algebra.

Let S_n^+ (resp. S_n^-) be the subalgebra of S_n generated by $H_{i,l}$ ($0 \leq i \leq n$, $l > 0$) (resp. $0 \leq i \leq n$, $l < 0$).

We introduce the Fock space

$$\mathcal{F}_n = S_n v_0$$

with the defining relations:

$$H_{i,l} v_0 = 0, \quad \text{for } l > 0, \quad (3.1.2)$$

$$q^{\frac{1}{2}c} v_0 = q^{\frac{1}{2}} v_0. \quad (3.1.3)$$

Note that \mathcal{F}_n is a free S_n^- -module of rank 1.

Let \mathbb{F} be a field of characteristic zero and let \mathfrak{a} be an associative \mathbb{F} -algebra generated by x_p, y_p ($p \in \mathbb{Z}_{>0}$), z and its inverse z^{-1} with the following relations:

$$[x_p, z] = [y_p, z] = 0,$$

$$\begin{aligned} [x_p, x_r] &= [y_p, y_r] = 0, \\ [x_p, y_r] &= \delta_{pr} z. \end{aligned}$$

Let \mathfrak{a}^+ (*resp.* \mathfrak{a}^-) be the subalgebra of \mathfrak{a} generated by x_p (*resp.* y_p). We set $\mathfrak{b} = \mathfrak{a}^+ \otimes \mathbb{F}[z, z^{-1}]$. This is a maximal abelian subalgebra of \mathfrak{a} . Fix a nonzero scalar $\lambda \in \mathbb{F}^\times$. Let \mathbb{F}_λ be the one-dimensional space \mathbb{F} viewed as a \mathfrak{b} -module by:

$$z \cdot 1 = \lambda, \quad \mathfrak{a}^+ \cdot 1 = 0.$$

Let $F(\lambda)$ be the induced \mathfrak{a} -module

$$F(\lambda) = \text{Ind}_{\mathfrak{b}}^{\mathfrak{a}} \mathbb{F}_\lambda = \mathfrak{a} \otimes_{\mathfrak{b}} \mathbb{F}_\lambda.$$

By the defining relations of \mathfrak{a} we obtain an \mathbb{F} -linear isomorphism

$$F(\lambda) \cong \mathfrak{a}^-.$$

Since \mathfrak{a}^- is abelian we may regard it as the algebra of polynomials in the variables y_1, y_2, \dots . Then we see that z acts on $\mathfrak{a}^- \cong \mathbb{F}[y_1, y_2, \dots]$ by the multiplication of λ , x_p acts by $\lambda \frac{\partial}{\partial y_p}$. By this realization we immediately have the following lemma.

Lemma 3.1.1. *$F(\lambda)$ is an irreducible \mathfrak{a} -module.*

Fix an skew-symmetric $(n+1) \times (n+1)$ -matrix with integral coefficients $M = (m_{ij})_{0 \leq i, j \leq n}$. We say that $\kappa \in \mathbb{Q}(q)$ is generic with respect to M if for any $k \in \mathbb{Z}_{>0}$ the matrix $([k\langle h_i, \alpha_j \rangle] \kappa^{-km_{ij}})$ is invertible.

Note that if $n = 1$ any κ is not generic with respect to any M . Since the matrix $([k\langle h_i, \alpha_j \rangle])_{0 \leq i, j \leq n}$ is invertible for $n > 1$, there exists a generic κ for $n > 1$.

Lemma 3.1.2. (1) *For a fixed M , we assume that $\kappa \in \mathbb{Q}(q)$ is generic with respect to M (in particular $n > 1$). Then \mathcal{F}_n is an irreducible S_n -module.*

(2) *\mathcal{F}_1 is not irreducible.*

Proof. (1) Set $G(k) = (g(k)_{ij}) = ([k\langle h_i, \alpha_j \rangle] \kappa^{-km_{ij}})$. Since κ is generic with respect to M there exists its inverse $G(k)^{-1} = (g(k)^{ij})$ for any k . Note that by the definition $\sum_{0 \leq s \leq n} g(k)^{is} g(k)_{sj} = \delta_{ij}$.

We set

$$\tilde{H}_{i,k} = \begin{cases} \sum_{0 \leq s \leq n} \frac{k}{[k]} g(k)^{is} H_{s,k}, & \text{for } k > 0, \\ H_{i,k}, & \text{for } k < 0. \end{cases}$$

Then we have

$$\begin{aligned} [\tilde{H}_{i,k}, \tilde{H}_{j,l}] &= [\tilde{H}_{i,-k}, \tilde{H}_{j,-l}] = 0, \\ [\tilde{H}_{i,k}, \tilde{H}_{j,-l}] &= \delta_{k,l} \delta_{ij}, \end{aligned}$$

for $k, l > 0$. Since all $G(k)$ are regular, $\tilde{H}_{i,k}$ ($0 \leq i \leq n$, $k > 0$) generate S_n^+ .

We shall use Lemma 3.1.1. Put $\mathbb{F} = \mathbb{Q}(q)$, $\mathfrak{a} = \mathfrak{S}_n$, $\mathfrak{a}^\pm = S_n^\pm$, $\lambda = 1$, $x_p = \tilde{H}_{i,k}$ ($k > 0$), $y_r = \tilde{H}_{j,l}$ ($l < 0$) where $p = (k-1)(n+1) + i + 1$ and $r = (-l-1)(n+1) + j + 1$.

Then it is clear that $F(1) = \mathcal{F}_n$. By Lemma 3.1.1 we conclude that \mathcal{F}_n is an irreducible S_n -module.

(2) It is easy to see that $\kappa^{-km_{01}}H_{0,-k} + H_{1,-k}$ is a central element of S_1 for each $k \in \mathbb{Z}_{>0}$. Therefore \mathcal{F}_1 has infinitely many singular vectors. \square

3.2. Construction of level (1,0) modules. We assume $n > 1$ and M is an $(n+1) \times (n+1)$ -matrix defined as follows;

$$M = \begin{pmatrix} 0 & -a & 0 & \dots & 0 & a \\ a & 0 & -a & \dots & 0 & 0 \\ 0 & a & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & -a \\ -a & 0 & 0 & \dots & a & 0 \end{pmatrix}$$

here $a \in \mathbb{Z}$.

Note that one can rewrite $\bar{P} = \oplus_{i=2}^n \mathbb{Z}\bar{\alpha}_i \oplus \mathbb{Z}\bar{\Lambda}_n$. We introduce a twisted version of the group algebra $\mathbb{Q}(q)[\bar{P}]$ by $\mathbb{Z}/2\mathbb{Z}$. We denote it by $\mathbb{Q}(q)\{\bar{P}\}$. This is the $\mathbb{Q}(q)$ -algebra generated by symbols $e^{\bar{\alpha}_2}$, $e^{\bar{\alpha}_3}, \dots, e^{\bar{\alpha}_n}$, $e^{\bar{\Lambda}_n}$ which satisfy the following relations:

$$e^{\bar{\alpha}_i} e^{\bar{\alpha}_j} = (-1)^{\langle \bar{h}_i, \bar{\alpha}_j \rangle} e^{\bar{\alpha}_j} e^{\bar{\alpha}_i} \quad (3.2.1)$$

$$e^{\bar{\alpha}_i} e^{\bar{\Lambda}_n} = (-1)^{\delta_{in}} e^{\bar{\Lambda}_n} e^{\bar{\alpha}_i}. \quad (3.2.2)$$

For $\bar{\alpha} = \sum_{i=2}^n m_i \bar{\alpha}_i + m_{n+1} \bar{\Lambda}_n$ we denote $e^{\bar{\alpha}} = (e^{\bar{\alpha}_2})^{m_2} (e^{\bar{\alpha}_3})^{m_3} \dots (e^{\bar{\alpha}_n})^{m_n} (e^{\bar{\Lambda}_n})^{m_{n+1}}$. For example $e^{\bar{\alpha}_1} = e^{-2\bar{\alpha}_2} e^{-3\bar{\alpha}_3} \dots e^{-n\bar{\alpha}_n} e^{(n+1)\bar{\Lambda}_n}$, $e^{\bar{\Lambda}_i} = e^{-\bar{\alpha}_{i+1}} e^{-2\bar{\alpha}_{i+2}} \dots e^{-(n-i)\bar{\alpha}_n} e^{(n-i+1)\bar{\Lambda}_n}$ where $\bar{\Lambda}_i$ is the i -th fundamental weight. We denote $\bar{\alpha}_0 = -\sum_{i=1}^n \bar{\alpha}_i$ and $\bar{h}_0 = -\sum_{i=1}^n \bar{h}_i$.

Note that $\langle h_i, \alpha_j \rangle = \langle \bar{h}_i, \bar{\alpha}_j \rangle$ for $0 \leq i, j \leq n$.

We denote by $\mathbb{Q}(q)\{\bar{Q}\}$ the subalgebra of $\mathbb{Q}(q)\{\bar{P}\}$ generated by $e^{\bar{\alpha}_i}$ ($1 \leq i \leq n$).

Set

$$W(p)_n = \mathcal{F}_n \otimes \mathbb{Q}(q)\{\overline{Q}\}e^{\overline{\Lambda}_p} \quad \text{for } 0 \leq p \leq n.$$

We define the operators $H_{i,l}$, $e^{\overline{\alpha}}$ ($\overline{\alpha} \in \overline{Q}$), $\partial_{\overline{\alpha}_i}$, $z^{H_{i,0}}$, d on $W(p)_n$ for $i = 0, 1, \dots, n$ as follows:

for $v \otimes e^{\overline{\beta}} = H_{i_1, -k_1} \cdots H_{i_N, -k_N} v_0 \otimes e^{\sum_{j=1}^n m_j \overline{\alpha}_j + \overline{\Lambda}_p} \in W(p)_n$,

$$\begin{aligned} H_{i,l}(v \otimes e^{\overline{\beta}}) &= (H_{i,l}v) \otimes e^{\overline{\beta}}, \\ e^{\overline{\alpha}}(v \otimes e^{\overline{\beta}}) &= v \otimes e^{\overline{\alpha} + \overline{\beta}}, \\ \partial_{\overline{\alpha}_i}(v \otimes e^{\overline{\beta}}) &= \langle \overline{h}_i, \overline{\beta} \rangle v \otimes e^{\overline{\beta}}, \\ z^{H_{i,0}}(v \otimes e^{\overline{\beta}}) &= z^{\langle \overline{h}_i, \overline{\beta} \rangle} \kappa^{\frac{1}{2} \sum_{j=1}^n \langle \overline{h}_i, m_j \overline{\alpha}_j \rangle} v \otimes e^{\overline{\beta}}, \\ d(v \otimes e^{\overline{\beta}}) &= \left(- \sum_{s=1}^N k_s - \frac{(\overline{\beta}|\overline{\beta})}{2} + \frac{(\overline{\Lambda}_p|\overline{\Lambda}_p)}{2} \right) v \otimes e^{\overline{\beta}}. \end{aligned}$$

We have the following lemma.

Lemma 3.2.1. *As operators on $W(p)_n$,*

$$\begin{aligned} e^{\overline{\alpha}_i} e^{\overline{\alpha}_j} &= (-1)^{\langle \overline{h}_i, \overline{\alpha}_j \rangle} e^{\overline{\alpha}_j} e^{\overline{\alpha}_i}, \\ e^{\overline{\alpha}_i} q^{\partial_{\overline{\alpha}_j}} &= q^{\langle \overline{h}_i, \overline{\alpha}_j \rangle} q^{\partial_{\overline{\alpha}_j}} e^{\overline{\alpha}_i}, \\ e^{\overline{\alpha}_i} z^{\partial_{\overline{\alpha}_j}} &= z^{\langle \overline{h}_i, \overline{\alpha}_j \rangle} z^{\partial_{\overline{\alpha}_j}} e^{\overline{\alpha}_i} \end{aligned}$$

for $0 \leq i, j \leq n$.

We introduce the following generating functions:

$$\begin{aligned} E_i(z) &= \sum_{k \in \mathbb{Z}} E_{i,k} z^{-k}, \\ F_i(z) &= \sum_{k \in \mathbb{Z}} F_{i,k} z^{-k}. \end{aligned}$$

Proposition 3.2.2. *Assume $c = 1$ and $n > 1$. Then for each p and κ , the following action gives a $U_q^{\phi'}(\mathfrak{g}_{\text{tor}})$ -module structure on $W(p)_n$:*

$$\begin{aligned} q^{\frac{1}{2}c} &\mapsto q^{\frac{1}{2}}, \\ q^{d_1} &\mapsto q^d, \\ E_i(z) &\mapsto \exp\left(\sum_{k \geq 1} \frac{H_{i,-k}}{[k]} (q^{-1/2}z)^k\right) \exp\left(\sum_{k \geq 1} -\frac{H_{i,k}}{[k]} (q^{1/2}z)^{-k}\right) e^{\overline{\alpha}_i} z^{H_{i,0}+1}, \\ F_i(z) &\mapsto \exp\left(\sum_{k \geq 1} -\frac{H_{i,-k}}{[k]} (q^{1/2}z)^k\right) \exp\left(\sum_{k \geq 1} \frac{H_{i,k}}{[k]} (q^{-1/2}z)^{-k}\right) e^{-\overline{\alpha}_i} z^{-H_{i,0}+1}, \\ K_i^+(z) &\mapsto \exp((q - q^{-1}) \sum_{k \geq 1} H_{i,k} z^{-k}) q^{\partial_{\overline{\alpha}_i}}, \\ K_i^-(z) &\mapsto \exp(-(q - q^{-1}) \sum_{k \geq 1} H_{i,-k} z^k) q^{-\partial_{\overline{\alpha}_i}} \end{aligned}$$

for $0 \leq i \leq n$.

The proof will be given in Appendix.

We have immediately the following lemma.

Lemma 3.2.3. *The $U_q^{\phi'}(\mathfrak{g}_{tor})$ -module $W(p)_n$ is cyclic: $W(p)_n = U_q^{\phi'}(\mathfrak{g}_{tor})(v_0 \otimes e^{\overline{\Lambda}_p})$.*

Theorem 3.2.4. *If $n > 1$ and κ is generic then $W(p)_n$ is irreducible for any p .*

Proof. Since \mathcal{F}_n is irreducible with respect to the action of S_n , it is enough to show that for any non-zero $v = v_0 \otimes \sum_{\overline{\alpha} \in \overline{Q}} a_{\overline{\alpha}} e^{\overline{\alpha}} e^{\overline{\Lambda}_i}$ ($a_{\overline{\alpha}} \in \mathbb{Q}(q)$) there exists $X \in U_q^{\phi'}(\mathfrak{g}_{tor})$ such that $Xv = v_0 \otimes e^{\overline{\Lambda}_p}$. Let \overline{S}_n be the subalgebra of S_n generated by $H_{i,l}$ ($1 \leq i \leq n, l \in \mathbb{Z} \setminus \{0\}$) and $q^{\frac{1}{2}c}$. Let $\overline{\mathcal{F}}_n$ be the \overline{S}_n -submodule of \mathcal{F}_n generated by $v_0 \otimes e^{\overline{\Lambda}_i}$, and let $\overline{W(p)}_n = \overline{\mathcal{F}}_n \otimes \mathbb{Q}(q)\{\overline{Q}\}e^{\overline{\Lambda}_p}$. As already known $\overline{W(p)}_n$ is an irreducible $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module. It is obvious that $v \in \overline{W(p)}_n$. Therefore there exists $X \in U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1}) \subset U_q^{\phi'}(\mathfrak{g}_{tor})$ such that $Xv = v_0 \otimes e^{\overline{\Lambda}_p}$. \square

Remark 3.2.5. Since $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ and $U_q^{(2)'}(\widehat{\mathfrak{sl}}_{n+1})$ are subalgebras of $U_q^{\phi'}(\mathfrak{g}_{tor})$ we can regard $W(p)_n$ as a $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module or as a $U_q^{(2)'}(\widehat{\mathfrak{sl}}_{n+1})$ -module. As a $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module, $W(p)_n$ is a level 1 module. On the other hand it is a level 0 module as a $U_q^{(2)'}(\widehat{\mathfrak{sl}}_{n+1})$ -module.

3.3. On the structure of level (1,0) modules. In this subsection we will study the $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module structure of $W(p)_n$.

Let M be a $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module. We denote the character of M by ch_M .

Let $L(\Lambda_p)$ be the irreducible highest weight $U_q(\widehat{\mathfrak{sl}}_{n+1})$ -module with highest weight Λ_p . Note that the following identity holds:

$$\text{ch}_{L(\Lambda_p)} = \frac{e^{\Lambda_p} \sum_{\alpha \in \overline{Q}} e^{\alpha - (\frac{1}{2}(\alpha|\alpha) - (\alpha|\overline{\Lambda}_p))\delta}}{\varphi(e^{-\delta})^n}. \quad (3.3.1)$$

Here $\varphi(x) = \prod_{k>0} (1 - x^k)$.

We denote δ_1 by the null root of $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$.

By the definition of $W(p)_n$ and (3.3.1) it is immediate to see the following proposition.

Proposition 3.3.1.

$$\begin{aligned} ch_{W(p)_n} &= \frac{e^{\Lambda_p} \sum_{\alpha \in \overline{\mathbb{Q}}} e^{\alpha - (\frac{1}{2}(\alpha|\alpha) - (\alpha|\overline{\Lambda_p}))\delta_1}}{\varphi(e^{-\delta})^n} \\ &= \frac{ch_{L(\Lambda_p)}}{\varphi(e^{-\delta_1})}. \end{aligned}$$

Lemma 3.3.2. *For each $l \in \mathbb{Z} \setminus \{0\}$ there exist $\widehat{H}_l = \sum_{i=0}^n a_{i,l} H_{i,l}$ ($a_{i,l} \in \mathbb{Q}(q)$) such that*

$$[\widehat{H}_l, H_{j,k}] = 0$$

for any $1 \leq j \leq n$ and $k \in \mathbb{Z} \setminus \{0\}$. Moreover such \widehat{H}_l is unique up to scalar.

Proof. Note that $\kappa = 1$. The rank of $n \times (n+1)$ -matrix $([l\langle h_i, \alpha_j \rangle])_{1 \leq i \leq n, 0 \leq j \leq n}$ is equal to n . The lemma follows from this fact immediately. \square

By the definition of \widehat{H}_l we have

$$[\widehat{H}_k, \widehat{H}_l] = \delta_{k+l,0} \gamma_k, \quad (3.3.2)$$

where $\gamma_k \in \mathbb{Q}(q)$. We fix a normalization of \widehat{H}_l by putting $\gamma_k = 1$ for all k .

Let \widehat{S}_n be the subalgebra of S_n generated by \widehat{H}_l . By the definition, \widehat{S}_n acts on $W(p)_n$.

The following two lemmas are easy to see.

Lemma 3.3.3. *For $l > 0$, $\widehat{H}_l(v_0 \otimes e^{\bar{\beta}}) = 0$.*

Lemma 3.3.4. *The action of $U_q^{(1')}(\widehat{\mathfrak{sl}}_{n+1})$ on $W(p)_n$ commutes with the action of \widehat{S}_n .*

Let \widehat{S}_n^- be the subalgebra of \widehat{S}_n generated by \widehat{H}_l ($l < 0$).

Proposition 3.3.5. *As $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module*

$$W(p)_n \cong L(\Lambda_p)^{\oplus \infty}.$$

Proof. Set $\deg(\widehat{H}_k) = k$. Let $M_k = M_k(\widehat{H}_{-1}, \widehat{H}_{-2}, \dots)$ be a monomial of degree k in variables $\widehat{H}_{-1}, \widehat{H}_{-2}, \dots$. Then by the above two lemmas, $M_k v_0 \otimes e^{\overline{\Lambda_p}}$ is a singular vector of $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module $W(p)_n$. Let W_{M_k} be the $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -submodule which is generated by $M_k v_0 \otimes e^{\overline{\Lambda_p}}$. Then by the definition of the action of $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ on $W(p)_n$, we have

$$W_{M_k} \cong L(\Lambda_p - k\delta) \cong L(\Lambda_p).$$

The vectors $\{M_k v_0 \otimes e^{\overline{\Lambda_p}}\}$ are linearly independent. The number of the monomials of degree k is equal to the k -th partition number $p(k)$. Therefore there is a $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -submodule W of $W(p)_n$ which is isomorphic to $\bigoplus_{k \geq 0} L(\Lambda_p - k\delta)^{\oplus p(k)}$. By Proposition 3.3.1 it coincides with $W(p)_n$. This completes proof. \square

By Lemma 3.3.4 and the proof of Proposition 3.3.5, the following corollary follows immediately.

Corollary 3.3.6. *As $U_q^{(1)' }(\widehat{\mathfrak{sl}}_{n+1}) \otimes \widehat{S}_n$ -module $W(p)_n$ is isomorphic to $L(\Lambda_p) \otimes \widehat{S}_n^-$.*

3.4. Construction of level (1,1) modules. We introduce a twisted version of the group algebra $\mathbb{Q}(q)[Q]$ by $\mathbb{Z} \setminus 2\mathbb{Z}$. We denoted it by $\mathbb{Q}(q)\{Q\}$. This is the $\mathbb{Q}(q)$ -algebra generated by symbols $e^{\alpha_0}, e^{\alpha_1}, \dots, e^{\alpha_n}$ which satisfy the following relations:

$$e^{\alpha_i} e^{\alpha_j} = (-1)^{\langle h_i, \alpha_j \rangle} e^{\alpha_j} e^{\alpha_i}. \quad (3.4.1)$$

Similarly to §3.2, we denote $e^\alpha = (e^{\alpha_0})^{m_0} (e^{\alpha_1})^{m_1} \dots (e^{\alpha_n})^{m_n}$ for $\alpha = \sum_{i=0}^n m_i \alpha_i \in Q$.

Let

$$V(p)_n = \mathcal{F}_n \otimes \mathbb{Q}(q)\{Q\} e^{\Lambda_p}.$$

Here we regard e^{Λ_p} only a symbol indexed by p .

We define the operators $H_{i,l}$ ($0 \leq i \leq n$, $l \neq 0$), e^α ($\alpha \in Q$), ∂_{α_i} and $z^{H_{i,0}}$ ($0 \leq i \leq n$) on $V(p)_n$ as follows:

for $v \otimes e^\beta e^{\Lambda_p} = H_{i_1, -k_1} \dots H_{i_N, -k_N} v_0 \otimes e^\beta e^{\Lambda_p} \in V(p)_n$ ($\beta = \sum_{k=0}^n m_k \alpha_k \in Q$),

$$\begin{aligned} H_{i,l}(v \otimes e^\beta e^{\Lambda_p}) &= (H_{i,l}v) \otimes e^\beta e^{\Lambda_p}, \\ e^\alpha(v \otimes e^\beta e^{\Lambda_p}) &= v \otimes (e^\alpha e^\beta) e^{\Lambda_p}, \\ \partial_{\alpha_i}(v \otimes e^\beta e^{\Lambda_p}) &= \langle h_i, \beta + \Lambda_p \rangle v \otimes e^\beta e^{\Lambda_p}, \\ z^{H_{i,0}}(v \otimes e^\beta e^{\Lambda_p}) &= z^{\langle h_i, \beta + \Lambda_p \rangle} \kappa^{\frac{1}{2} \sum_{k=0}^n \langle h_i, m_k \alpha_k \rangle m_{ik}} v \otimes e^\beta e^{\Lambda_p}, \\ d_1(v \otimes e^\beta e^{\Lambda_p}) &= \left(- \sum_{s=1}^N k_s - \frac{(\beta|\beta)}{2} - (\beta|\Lambda_p) \right) v \otimes e^\beta e^{\Lambda_p}, \\ d_2(v \otimes e^\beta e^{\Lambda_p}) &= m_0(v \otimes e^\beta e^{\Lambda_p}). \end{aligned}$$

The following lemma is easy.

Lemma 3.4.1. *As operators on $V(p)_n$,*

$$\begin{aligned} e^{\alpha_i} e^{\alpha_j} &= (-1)^{\langle h_i, \alpha_j \rangle} e^{\alpha_j} e^{\alpha_i} \\ q^{\partial_{\alpha_i}} e^{\alpha_j} &= q^{\langle h_i, \alpha_j \rangle} e^{\alpha_j} q^{\partial_{\alpha_i}} \\ z^{H_{i,0}} e^{\alpha_j} &= z^{\langle h_i, \alpha_j \rangle} \kappa^{\frac{1}{2} \langle h_i, \alpha_j \rangle m_{ij}} e^{\alpha_j} z^{H_{i,0}}. \end{aligned}$$

Proposition 3.4.2. *Assume $c = 1$ and $n > 1$. Then for each p , the following action gives a $U_q(\mathfrak{g}_{tor})$ -module structure on $V(p)_n$:*

$$\begin{aligned} q^{\frac{1}{2}c} &\mapsto q^{\frac{1}{2}}, \\ q^{d_1} &\mapsto q^{d_1} \\ q^{d_2} &\mapsto q^{d_2} \\ E_i(z) &\mapsto \exp\left(\sum_{k \geq 1} \frac{H_{i,-k}}{[k]} (q^{-1/2}z)^k\right) \exp\left(\sum_{k \geq 1} -\frac{H_{i,k}}{[k]} (q^{1/2}z)^{-k}\right) e^{\alpha_i} z^{H_{i,0}+1}, \\ F_i(z) &\mapsto \exp\left(\sum_{k \geq 1} -\frac{H_{i,-k}}{[k]} (q^{1/2}z)^k\right) \exp\left(\sum_{k \geq 1} \frac{H_{i,k}}{[k]} (q^{-1/2}z)^{-k}\right) e^{-\alpha_i} z^{-H_{i,0}+1}, \\ K_i^+(z) &\mapsto \exp\left((q - q^{-1}) \sum_{k \geq 1} H_{i,k} z^{-k}\right) q^{\partial_{\alpha_i}}, \\ K_i^-(z) &\mapsto \exp\left(-(q - q^{-1}) \sum_{k \geq 1} H_{i,-k} z^k\right) q^{-\partial_{\alpha_i}} \end{aligned}$$

for $0 \leq i \leq n$.

The proof will be given in Appendix.

It is easy to see the following lemma.

Lemma 3.4.3. *$V(p)_n$ is a cyclic $U_q(\mathfrak{g}_{tor})$ -module: $V(p)_n = U_q(\mathfrak{g}_{tor})(v_0 \otimes e^{\Lambda_p})$.*

Lemma 3.4.4. *$V(p)_n$ has level 1 as a $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module and as a $U_q^{(2)}(\widehat{\mathfrak{sl}}_{n+1})$ -module.*

Proof. It is clear that $V(p)_n$ is a level 1 $U_q^{(1)}(\widehat{\mathfrak{sl}}_{n+1})$ -module. The center of $U_q^{(2)}(\widehat{\mathfrak{sl}}_{n+1})$ is $\prod_{k=0}^n K_i$. By the definition it acts as the scalar q on $V(p)_n$. \square

4. ON $U_q(\mathfrak{sl}_{2,tor})$

4.1. In this section we assume that $\mathfrak{g} = \mathfrak{sl}_2$. We shall try to find finitely many generators of $U_q(\mathfrak{sl}_{2,tor})$.

Let

$$\begin{aligned} E_i &= E_{i,0}, \quad F_i = F_{i,0}, \quad q^{\pm h_i} = K_i^{\pm}, \text{ for } i = 0, 1, \\ E_{-1} &= F_{0,1} K_0^-, \quad F_{-1} = K_0^+ E_{0,-1}, \quad q^{\pm h_{-1}} = q^{\pm c} K_0^{\mp}. \end{aligned}$$

Proposition 4.1.1. *$U_q(\mathfrak{sl}_{2,tor})$ is generated by $E_i, F_i, q^{\pm h_i}$ ($i = -1, 0, 1$), $q^{\pm \frac{1}{2}c}, q^{\pm d_1}, q^{\pm d_2}$.*

Proof. Let \mathcal{A} be the subalgebra of $U_q(\mathfrak{sl}_{2,tor})$ generated by $E_i, F_i, q^{\pm h_i}$ ($i = -1, 0, 1$), $q^{\pm \frac{1}{2}c}$, $q^{\pm d_1}$, $q^{\pm d_2}$. By the definition we have $E_{0,-1} = q^{-h_0} F_{-1}$ and $F_{0,1} = E_{-1} q^{h_0}$. Since

$$\begin{aligned} [E_{0,0}, F_{0,1}] &= \frac{1}{q - q^{-1}} q^{-\frac{1}{2}c} K_{0,1}^+ \\ &= q^{-\frac{1}{2}c} q^{h_0} H_{0,1} \end{aligned}$$

and

$$\begin{aligned} [E_{0,-1}, F_{0,0}] &= -\frac{1}{q - q^{-1}} q^{\frac{1}{2}c} K_{0,-1}^- \\ &= q^{\frac{1}{2}c} q^{-h_0} H_{0,-1}. \end{aligned}$$

we deduce $H_{0,1}$ and $H_{0,-1} \in \mathcal{A}$. We recall (2.2.7)

$$\begin{aligned} [H_{i,k}, E_{j,l}] &= \frac{1}{k} [k \langle h_i, \alpha_j \rangle] q^{-\frac{1}{2}|k|c} \kappa^{-km_{ij}} E_{j,k+l}, \\ [H_{i,k}, F_{j,l}] &= -\frac{1}{k} [k \langle h_i, \alpha_j \rangle] q^{\frac{1}{2}|k|c} \kappa^{-km_{ij}} F_{j,k+l}. \end{aligned}$$

By these formulas we have $E_{i,k}, F_{i,k} \in \mathcal{A}$ for $i = 0, 1, k \in \mathbb{Z}$ inductively.

On the other hand we know

$$\begin{aligned} [E_{0,1}, F_{0,1}] &= \frac{1}{q - q^{-1}} q^{-\frac{1}{2}c} K_{0,2}^+ \\ &= \frac{1}{2} (q - q^{-1}) H_{0,1}^2 + H_{0,2}. \end{aligned}$$

Therefore we get $H_{0,2} \in \mathcal{A}$. Similarly we have $H_{i,l} \in \mathcal{A}$ for any i, l .

This completes the proof. \square

Lemma 4.1.2. *The following relations hold in $U_q(\mathfrak{sl}_{2,tor})$:*

$$[q^{\pm h_i}, q^{\pm h_j}] = 0, \quad (4.1.1)$$

$$[q^{\pm d_i}, q^{\pm h_j}] = 0, \quad (4.1.2)$$

$$q^{d_1} E_j q^{-d_1} = q^{\delta_{j,-1}} E_j, \quad (4.1.3)$$

$$q^{d_1} F_j q^{-d_1} = q^{-\delta_{j,-1}} F_j,$$

$$q^{d_2} E_j q^{-d_2} = q^{-1+\delta_{j,1}} E_j, \quad (4.1.4)$$

$$q^{d_2} F_j q^{-d_2} = q^{1-\delta_{j,1}} F_j,$$

$$q^{h_i} E_j q^{-h_i} = q^{a_{ij}} E_j, \quad (4.1.5)$$

$$q^{h_i} F_j q^{-h_i} = q^{-a_{ij}} F_j,$$

where

$$(a_{ij})_{-1 \leq i, j \leq 1} = \begin{pmatrix} 2 & -2 & 2 \\ -2 & 2 & -2 \\ 2 & -2 & 2 \end{pmatrix},$$

$$[E_i, F_j] = \delta_{ij} \frac{q^{h_i} - q^{-h_i}}{q - q^{-1}}, \quad \text{for } |i - j| \leq 1, \quad (4.1.6)$$

$$E_{-1}^3 F_1 - q^{-2} [3] E_{-1}^2 F_1 E_{-1} + q^{-4} [3] E_{-1} F_1 E_{-1}^2 - q^{-6} F_1 E_{-1}^3 = 0, \quad (4.1.7)$$

$$E_1^3 F_{-1} - q^2 [3] E_1^2 F_{-1} E_1 + q^4 [3] E_1 F_{-1} E_1^2 - q^6 F_{-1} E_1^3 = 0, \quad (4.1.8)$$

$$F_{-1}^3 E_1 - q^{-2} [3] F_{-1}^2 E_1 F_{-1} + q^{-4} [3] F_{-1} E_1 F_{-1}^2 - q^{-6} E_1 F_{-1}^3 = 0, \quad (4.1.9)$$

$$F_1^3 E_{-1} - q^2 [3] F_1^2 E_{-1} F_1 + q^4 [3] F_1 E_{-1} F_1^2 - q^6 E_{-1} F_1^3 = 0, \quad (4.1.10)$$

$$E_i^3 E_j - [3] E_i^2 E_j E_i + [3] E_i E_j E_i^2 - E_j E_i^3 = 0, \quad \text{for } |i - j| = 1, \quad (4.1.11)$$

$$F_i^3 F_j - [3] F_i^2 F_j F_i + [3] F_i F_j F_i^2 - F_j F_i^3 = 0, \quad \text{for } |i - j| = 1, \quad (4.1.12)$$

$$E_{-1} E_1 - q^2 E_1 E_{-1} = 0, \quad (4.1.13)$$

$$F_{-1} F_1 - q^2 F_1 F_{-1} = 0. \quad (4.1.14)$$

Proof. By the definition of $U_q(\mathfrak{g}_{tor})$ and E_i , F_i and q^{h_i} it is easy to check these relations. \square

Let \mathcal{U} be an associative algebra over $\mathbb{Q}(q)$ generated by E_i , F_i , $q^{\pm h_i}$ ($i = -1, 0, 1$), $q^{\pm \frac{1}{2}c}$, $q^{\pm d_1}$, $q^{\pm d_2}$ with relations (4.1.1)-(4.1.14). Then we have,

Corollary 4.1.3. *There is a canonical surjective algebra homomorphism $\Psi : \mathcal{U} \rightarrow U_q(\mathfrak{sl}_{2,tor})$.*

Remark 4.1.4. Ψ has a highly nontrivial kernel. It is important to decide it. For example the following formulas holds in $U_q(\mathfrak{sl}_{2,tor})$:

$$\kappa^{m_{01}} E_{0,0} E_{1,-1} - q^{-2} \kappa^{m_{01}} E_{1,-1} E_{0,0} = q^{-2} E_{0,-1} E_{1,0} - E_{1,0} E_{0,-1},$$

$$E_{0,-1} = q^{-h_0} F_{-1}$$

and

$$E_{1,-1} = \frac{\kappa^{-m_{01}}}{[-2]} [F_{-1} F_0 - q^{-2} F_0 F_{-1}, E_1].$$

Therefore we have

$$\begin{aligned} X &= \frac{1}{[-2]} E_0 [F_{-1} F_0 - q^{-2} F_0 F_{-1}, E_1] - \frac{q^{-2}}{[-2]} [F_{-1} F_0 - q^{-2} F_0 F_{-1}, E_1] E_0 \\ &\quad - q^{-2} q^{-h_0} F_{-1} E_1 - E_1 q^{-h_0} F_{-1} \\ &= 0 \end{aligned}$$

in $U_q(\mathfrak{sl}_{2,tor})$. Thus $X \in \text{Ker} \Psi$. But, as an element of \mathcal{U} , X is not equal to 0.

4.2. Let

$$E_{0*} = F_{1,1} K_1^-, \quad F_{0*} = K_1^+ E_{1,-1}, \quad q^{\pm h_{0*}} = q^{\pm c} K_1^{\mp}.$$

Proposition 4.2.1. *The subalgebra generated by E_i , F_i and q^{h_i} for $i = 0, 1$, E_{0*} , F_{0*} , $q^{\pm h_{0*}}$, $q^{\pm c}$, $q^{\pm d_1}$, $q^{\pm d_2}$ is equal to $U_q(\mathfrak{sl}_{2,tor})$. That is, they are generators of $U_q(\mathfrak{sl}_{2,tor})$. Moreover these generators satisfy relations similar to the ones in Lemma 4.1.2.*

Proof. This proposition is proved in the same way as Proposition 4.1.1 and Lemma 4.1.2. \square

We have immediately the following lemma.

Lemma 4.2.2. *Let $U_q^{(1)}(\widehat{\mathfrak{sl}}_2)$ be the subalgebras generated by E_i , F_i , q^{h_i} for $i = 1, 0^*$ and $q^{\pm(d_1+d_2)}$, $U_q^{(2)}(\widehat{\mathfrak{sl}}_2)$ the subalgebras generated by E_i , F_i , q^{h_i} for $i = 0, 1$ and $q^{\pm(d_1+d_2)}$, $U_q^{(3)}(\widehat{\mathfrak{sl}}_2)$ the subalgebras generated by E_i , F_i , $q^{\pm h_i}$ for $i = 0^*, -1$ and $q^{\pm(d_1+d_2)}$, and $U_q^{(4)}(\widehat{\mathfrak{sl}}_2)$ the subalgebras generated by E_i , F_i , $q^{\pm h_i}$ for $i = 0, -1$ and $q^{\pm(d_1+d_2)}$. Then $U_q^{(i)}$ ($i = 1, 2, 3, 4$) are isomorphic to $U_q(\widehat{\mathfrak{sl}}_2)$.*

Those four algebras are schematically visualized by Fig. 1.




Fig. 1.

Let $U_q(\mathfrak{sl}_2)_{(i)}$ ($i = -1, 0, 1, 0^*$) be the sub algebra of $U_q(\mathfrak{sl}_{2,\text{tot}})$ generated by $E_i, F_i, q^{\pm h_i}$. All $U_q(\mathfrak{sl}_{2,(i)})$ ($i \in \{-1, 0, 1, 0^*\}$) are isomorphic to $U_q(\mathfrak{sl}_2)$. The upper left circle in Fig. 1 means $U_q(\mathfrak{sl}_2)_{(0)}$, the upper right one means $U_q(\mathfrak{sl}_2)_{(1)}$, the lower left one means $U_q(\mathfrak{sl}_2)_{(-1)}$ and the lower right one means $U_q(\mathfrak{sl}_2)_{(0^*)}$. The diagram

$$\circ^i \Longleftrightarrow \circ^j \quad (i, j \in \{-1, 0, 1, 0^*\})$$

means the algebra generated by $U_q(\mathfrak{sl}_2)_{(i)}$ and $U_q(\mathfrak{sl}_2)_{(j)}$ is isomorphic to $U_q(\widehat{\mathfrak{sl}}_2)$. For example $\circ^0 \Longleftrightarrow \circ^1$ means the algebra generated by $U_q(\mathfrak{sl}_2)_{(0)}$ and $U_q(\mathfrak{sl}_2)_{(1)}$ which we call $U_q^{(2)}(\widehat{\mathfrak{sl}}_2)$ is isomorphic to $U_q(\widehat{\mathfrak{sl}}_2)$. The meaning of the diagram

$$\circ^i \Longrightarrow \circ^j$$

is as follows: In the algebra generated by $U_q(\mathfrak{sl}_2)_{(i)}$ and $U_q(\mathfrak{sl}_2)_{(j)}$, the following relations hold

$$q^{h_i} E_j q^{-h_i} = q^2 E_j, \quad q^{h_j} E_i q^{-h_j} = q^2 E_i, \quad (4.2.1)$$

$$q^{h_i} F_j q^{-h_i} = q^{-2} F_j, \quad q^{h_j} F_i q^{-h_j} = q^{-2} F_i, \quad (4.2.2)$$

$$E_i^3 F_j - q^{-2}[3] E_i^2 F_j E_i + q^{-4}[3] E_i F_j E_i^2 - q^{-6} F_j E_i^3 = 0, \quad (4.2.3)$$

$$\begin{aligned} E_j^3 F_i - q^2[3] E_j^2 F_i E_j + q^4[3] E_j F_i E_j^2 - q^6 F_i E_j^3 &= 0, \\ F_i^3 E_j - q^{-2}[3] F_i^2 E_j F_i + q^{-4}[3] F_i E_j F_i^2 - q^{-6} E_j F_i^3 &= 0, \end{aligned} \quad (4.2.4)$$

$$F_j^3 E_i - q^2[3] F_j^2 E_i F_j + q^4[3] F_j E_i F_j^2 - q^6 E_i F_j^3 = 0,$$

$$E_i E_j - q^2 E_j E_i = 0, \quad (4.2.5)$$

$$F_i F_j - q^2 F_j F_i = 0. \quad (4.2.6)$$

APPENDIX A.

A.1. Proof of Proposition 3.2.2 and 3.4.2. For the proof we rewrite the defining relation of $U_q(\mathfrak{g}_{tor})$ generating function level.

$$q^{\pm \frac{1}{2}c} \text{ are central,} \quad (\text{A.1.1})$$

$$K_i^+ K_i^- = K_i^- K_i^+ = 1, \quad (\text{A.1.2})$$

$$K_i^\pm(z) K_j^\pm(w) = K_j^\pm(w) K_i^\pm(z) \quad (\text{A.1.3})$$

$$\theta_{-\langle h_i, \alpha_j \rangle} (q^{-c} \kappa^{m_{ij}} \frac{z}{w}) K_i^-(z) K_j^+(w) = \theta_{-\langle h_i, \alpha_j \rangle} (q^c \kappa^{m_{ij}} \frac{z}{w}) K_j^+(w) K_i^-(z) \quad (\text{A.1.4})$$

$$q^{d_1} K_j^\pm(z) q^{-d_1} = K_j^\pm(q^{-1}z), \quad (\text{A.1.5})$$

$$[q^{d_2}, K_j^\pm(z)] = 0, \quad (\text{A.1.6})$$

$$q^{d_1} E_j(z) q^{-d_1} = E_j(q^{-1}z), \quad (\text{A.1.7})$$

$$q^{d_1} F_j(z) q^{-d_1} = F_j(q^{-1}z),$$

$$q^{d_2} E_j(z) q^{-d_2} = q^{\delta_{j0}} E_j(z), \quad (\text{A.1.8})$$

$$q^{d_2} F_j(z) q^{-d_2} = q^{-\delta_{j0}} F_j(z),$$

$$K_i^+(z) E_j(w) = \theta_{-\langle h_i, \alpha_j \rangle} (q^{-\frac{1}{2}c} \kappa^{-m_{ij}} \frac{w}{z}) E_j(w) K_i^+(z) \quad (\text{A.1.9})$$

$$K_i^-(z) E_j(w) = \theta_{\langle h_i, \alpha_j \rangle} (q^{-\frac{1}{2}c} \kappa^{m_{ij}} \frac{z}{w}) E_j(w) K_i^-(z)$$

$$K_i^+(z) F_j(w) = \theta_{\langle h_i, \alpha_j \rangle} (q^{\frac{1}{2}c} \kappa^{-m_{ij}} \frac{w}{z}) F_j(w) K_i^+(z)$$

$$K_i^-(z) F_j(w) = \theta_{-\langle h_i, \alpha_j \rangle} (q^{\frac{1}{2}c} \kappa^{m_{ij}} \frac{z}{w}) F_j(w) K_i^-(z)$$

$$[E_i(z), F_j(w)] = \delta_{i,j} \frac{1}{q - q^{-1}} \left\{ \delta(q^c \frac{w}{z}) K_i^+(q^{\frac{1}{2}c} w) - \delta(q^c \frac{z}{w}) K_i^-(q^{\frac{1}{2}c} z) \right\} \quad (\text{A.1.10})$$

$$(\kappa^{m_{ij}} z - q^{\langle h_i, \alpha_j \rangle} w) E_i(z) E_j(w) = (q^{\langle h_i, \alpha_j \rangle} \kappa^{m_{ij}} z - w) E_j(w) E_i(z) \quad (\text{A.1.11})$$

$$(\kappa^{m_{ij}} z - q^{-\langle h_i, \alpha_j \rangle} w) F_i(z) F_j(w) = (q^{-\langle h_i, \alpha_j \rangle} \kappa^{m_{ij}} z - w) F_j(w) F_i(z)$$

$$\sum_{\sigma \in \mathfrak{S}_m} \sum_{r=0}^m (-1)^r \begin{bmatrix} m \\ r \end{bmatrix}_q E_i(z_{\sigma(1)}) \cdots E_i(z_{\sigma(r)}) E_j(w) E_i(z_{\sigma(r+1)}) \cdots E_i(z_{\sigma(m)}) = 0 \quad (\text{A.1.12})$$

$$\sum_{\sigma \in \mathfrak{S}_m} \sum_{r=0}^m (-1)^r \begin{bmatrix} m \\ r \end{bmatrix}_q F_i(z_{\sigma(1)}) \cdots F_i(z_{\sigma(r)}) F_j(w) F_i(z_{\sigma(r+1)}) \cdots F_i(z_{\sigma(m)})$$

where $i \neq j$ and $m = 1 - \langle h_i, \alpha_j \rangle$.

In these formulas we denote $\theta_m(z) = \frac{zq^m - 1}{z - q^m}$ for $m \in \mathbb{Z}$, $\delta(z) = \sum_{k \in \mathbb{Z}} z^k$.

If Proposition 3.4.2 holds, then, from Lemma 3.2.1 and 3.4.1, we have Proposition 3.2.2 by putting $h_i \mapsto \bar{h}_i$, $\alpha_i \mapsto \bar{\alpha}_i$, $\kappa \mapsto 1$ and $z^{H_{i,0}} \mapsto z^{\bar{\delta}_i}$. Therefore it is enough to show Proposition 3.4.2.

The relations (A.1.1), (A.1.2) and (A.1.3) are trivial. (A.1.4) is just the commutation relations of Heisenberg algebra S_n . Therefore, by the definition of $V(p)_n$, it is clear that (A.1.4) holds. The relations (A.1.5), (A.1.6) immediately follows from the definition of d_1 and d_2 .

Let us show (A.1.7) and (A.1.8). Take $v \otimes e^\beta e^{\Lambda_p} \in V(p)_n$ where $\beta = \sum_{k=0}^n m_k \alpha_k \in Q$. Then we have

$$\begin{aligned} & q^{d_1} e^{\pm \alpha_j} z^{\pm H_{j,0}+1} q^{-d_1} (v \otimes e^\beta e^{\Lambda_p}) \\ &= q^{\mp(\alpha_j | \beta + \Lambda_p) - 1} z^{\pm \langle h_j, \beta + \Lambda_p \rangle + 1} \kappa^{\frac{1}{2} \sum_{k=0}^n \langle h_j, m_k \alpha_k \rangle m_{jk}} v \otimes e^{\pm \alpha_j} e^\beta e^{\Lambda_p} \\ &= (q^{-1} z)^{\pm \langle h_j, \beta + \Lambda_p \rangle + 1} \kappa^{\frac{1}{2} \sum_{k=0}^n \langle h_j, m_k \alpha_k \rangle m_{jk}} v \otimes e^{\pm \alpha_j} e^\beta e^{\Lambda_p} \\ &= e^{\pm \alpha_j} (q^{-1} z)^{\pm H_{j,0}+1} (v \otimes e^\beta e^{\Lambda_p}). \end{aligned}$$

Therefore we have

$$\begin{aligned} q^{d_1} E_j(z) q^{-d_1} &= q^{d_1} \exp\left(\sum_{k \geq 1} \frac{H_{j,-k}}{[k]} q^{-\frac{1}{2}k} z^k\right) \exp\left(-\sum_{k \geq 1} \frac{H_{j,k}}{[k]} q^{-\frac{1}{2}k} z^{-k}\right) e^{\alpha_j} z^{H_{j,0}+1} q^{-d_1} \\ &= \exp\left(\sum_{k \geq 1} \frac{H_{j,-k}}{[k]} q^{-\frac{1}{2}k} (q^{-1} z)^k\right) \exp\left(-\sum_{k \geq 1} \frac{H_{j,k}}{[k]} q^{-\frac{1}{2}k} (q^{-1} z)^{-k}\right) e^{\alpha_j} (q^{-1} z)^{H_{j,0}+1} \\ &= E_j(q^{-1} z). \end{aligned}$$

Similarly we have $q^{d_1} F_j(z) q^{-d_1} = F_j(q^{-1} z)$.

It is clear that

$$q^{d_2} e^{\pm \alpha_j} z^{\pm H_{j,0}+1} q^{-d_2} (v \otimes e^\beta e^{\Lambda_p}) = q^{\pm \delta_{j0}} e^{\pm \alpha_j} z^{\pm H_{j,0}+1} (v \otimes e^\beta e^{\Lambda_p}). \quad (\text{A.1.13})$$

From (A.1.13) and the fact that q^{d_2} commutes with $H_{j,k}$, we have (A.1.8).

We shall show (A.1.9). We denote

$$\begin{aligned} E_i^+(z) &= \exp\left(\sum_{k \geq 1} \frac{H_{i,-k}}{[k]} q^{-\frac{1}{2}k} z^k\right), \\ E_i^-(z) &= \exp\left(-\sum_{k \geq 1} \frac{H_{i,k}}{[k]} q^{-\frac{1}{2}k} z^{-k}\right), \\ F_i^+(z) &= \exp\left(-\sum_{k \geq 1} \frac{H_{i,-k}}{[k]} q^{\frac{1}{2}k} z^k\right), \\ F_i^-(z) &= \exp\left(\sum_{k \geq 1} \frac{H_{i,k}}{[k]} q^{\frac{1}{2}k} w^{-k}\right). \end{aligned}$$

Let us proof

$$K_i^+(z) E_j(w) = \theta_{-\langle h_i, \alpha_j \rangle} (q^{-\frac{1}{2}} \kappa^{m_{-ij}} \frac{w}{z}) E_j(w) K_i^+(z)$$

We have

$$\begin{aligned} [(q - q^{-1}) \sum_{k \geq 1} H_{i,k} z^{-k}, \sum_{l \geq 1} \frac{H_{j,-l}}{[l]} q^{-\frac{1}{2}l} w^l] &= \sum_{k,l} (q - q^{-1}) \frac{1}{[l]} [H_{i,k}, H_{j,-l}] z^{-k} q^{-\frac{1}{2}l} w^l \\ &= \sum_k (q - q^{-1}) \frac{[k \langle h_i, \alpha_j \rangle]}{k} \kappa^{-km_{ij}} q^{-\frac{1}{2}k} \left(\frac{w}{z}\right)^k \\ &= \sum_k \frac{1}{k} (q^{k(\langle h_i, \alpha_j \rangle - \frac{1}{2})} - q^{k(-\langle h_i, \alpha_j \rangle - \frac{1}{2})}) \kappa^{-km_{ij}} \left(\frac{w}{z}\right)^k \\ &= \log \frac{1 - q^{-\langle h_i, \alpha_j \rangle - \frac{1}{2}} \kappa^{-m_{ij}} \frac{w}{z}}{1 - q^{\langle h_i, \alpha_j \rangle - \frac{1}{2}} \kappa^{-m_{ij}} \frac{w}{z}} \end{aligned}$$

and

$$[(q - q^{-1}) \sum_{k \geq 1} H_{i,k} z^{-k}, -\sum_{l \geq 1} \frac{H_{j,l}}{[l]} q^{-\frac{1}{2}l} w^{-l}] = 0.$$

We recall Campbell-Hausdorff formula: let A and B be noncommutative operators and $C = [A, B]$. If $[C, A] = [C, B] = 0$ then we have $e^A e^B = e^C e^B e^A$.

By Campbell-Hausdorff formula we get

$$\begin{aligned} & \exp((q - q^{-1}) \sum_{k \geq 1} H_{i,k} z^{-k}) E_j^+(w) E_j^-(w) \\ &= \frac{1 - q^{-\langle h_i, \alpha_j \rangle - \frac{1}{2}} \kappa^{-m_{ij}} \frac{w}{z}}{1 - q^{\langle h_i, \alpha_j \rangle - \frac{1}{2}} \kappa^{-m_{ij}} \frac{w}{z}} E_j^+(w) E_j^-(w) \exp((q - q^{-1}) \sum_{k \geq 1} H_{i,k} z^{-k}). \end{aligned}$$

On the other hand, by Lemma 3.4.1 we have

$$q^{\partial_{\alpha_i}} e^{\alpha_j} w^{H_{j,0}} = q^{\langle h_i, \alpha_j \rangle} e^{\alpha_j} w^{H_{j,0}} q^{\partial_{\alpha_i}}.$$

Thus we get

$$\begin{aligned} & K_i^+(z) E_j(w) \\ &= \exp((q - q^{-1}) \sum_{k \geq 1} H_{i,k} z^{-k}) E_j^+(w) E_j^-(w) q^{\partial_{\alpha_i}} e^{\alpha_j} w^{H_{j,0}+1} \\ &= \frac{1 - q^{-\langle h_i, \alpha_j \rangle - \frac{1}{2}} \kappa^{-m_{ij}} \frac{w}{z}}{q^{-\langle h_i, \alpha_j \rangle - \frac{1}{2}} \kappa^{-m_{ij}} \frac{w}{z}} E_j(w) K_i^+(z) \\ &= \theta_{-\langle h_i, \alpha_j \rangle} (q^{-\frac{1}{2}} \kappa^{-m_{ij}} \frac{w}{z}) E_j(w) K_i^+(z). \end{aligned}$$

The other formulas in (A.1.9) can be checked by similar arguments.

Let us show (A.1.10). We have

$$\begin{aligned} & [- \sum_{k \geq 1} \frac{H_{i,k}}{[k]} q^{-\frac{1}{2}k} z^{-k}, - \sum_{k \geq 1} \frac{H_{j,-k}}{[k]} q^{\frac{1}{2}k} w^k] \\ &= \sum_{k \geq 1} \frac{1}{k[k]} [k \langle h_i, \alpha_j \rangle] \kappa^{-m_{ij}} \left(\frac{w}{z}\right)^k \\ &= \begin{cases} \log \frac{1}{(1-q\frac{w}{z})(1-q^{-1}\frac{w}{z})}, & \langle h_i, \alpha_j \rangle = 2 \quad (i = j), \\ \log(1 - \kappa^{-m_{ij}} \frac{w}{z}), & \langle h_i, \alpha_j \rangle = -1, \\ 0, & \langle h_i, \alpha_j \rangle = 0. \end{cases} \end{aligned}$$

For example we will show in the case of $\langle h_i, \alpha_j \rangle = -1$. By Campbell-Hausdorff formula we get

$$E_i^\pm(z) F_j^\pm(w) = F_j^\pm(w) E_i^\pm(z)$$

and

$$E_i^-(z) F_j^+(w) = (1 - \kappa^{-m_{ij}} \frac{w}{z}) F_j^+(w) E_i^-(z).$$

On the other hand, by Lemma 3.4.1, we have

$$z^{H_{i,0}} e^{-\alpha_j} = z \kappa^{\frac{1}{2}m_{ij}} e^{-\alpha_j} z^{H_{i,0}}.$$

Therefore we get

$$\begin{aligned} E_i(z)F_j(w) &= E_i^+(z)E_i^-(z)e^{\alpha_i}z^{H_{i,0}+1}F_j^+(w)F_j^-(w)e^{-\alpha_j}w^{-H_{j,0}+1} \\ &= (z\kappa^{\frac{1}{2}m_{ij}} - w\kappa^{-\frac{1}{2}m_{ij}})E_i^+(z)F_j^+(w)E_i^-(z)F_j^-(w) \\ &\quad \times e^{\alpha_i}e^{-\alpha_j}z^{H_{i,0}+1}w^{-H_{j,0}+1}. \end{aligned}$$

By a similar argument we have

$$\begin{aligned} F_j(w)E_i(z) &= -(w\kappa^{-\frac{1}{2}m_{ij}} - z\kappa^{\frac{1}{2}m_{ij}})E_i^+(z)F_j^+(w)E_i^-(z)F_j^-(w) \\ &\quad \times e^{\alpha_i}e^{-\alpha_j}z^{H_{i,0}+1}w^{-H_{j,0}+1}. \end{aligned}$$

Therefore we get

$$[E_i(z), F_j(w)] = 0.$$

Similarly one can check the other formulas.

We will show (A.1.11). We have

$$\begin{aligned} &[-\sum_{k \geq 1} \frac{H_{i,k}}{[k]} q^{-\frac{1}{2}k} z^{-k}, \sum_{k \geq 1} \frac{H_{j,-k}}{[k]} q^{-\frac{1}{2}k} w^k] \\ &= \sum_{k \geq 1} \frac{1}{k[k]} [k \langle h_i, \alpha_j \rangle] \kappa^{-m_{ij}} \left(\frac{w}{z}\right)^k \\ &= \begin{cases} \log(1 - \frac{w}{z})(1 - q^{-2}\frac{w}{z}), & i = j, \\ \log \frac{1}{1 - q^{-1}\kappa^{-m_{ij}}\frac{w}{z}}, & \langle h_i, \alpha_j \rangle = -1, \\ 0, & \langle h_i, \alpha_j \rangle = 0. \end{cases} \end{aligned}$$

For example let us show in the case of $\langle h_i, \alpha_j \rangle = -1$. If $\langle h_i, \alpha_j \rangle \neq -1$ one can show the formula by a similar argument. By Lemma 3.4.1 we get

$$\begin{aligned} &(z\kappa^{m_{ij}} - q^{\langle h_i, \alpha_j \rangle} w)E_i(z)E_j(w) \\ &= (z\kappa^{m_{ij}} - q^{-1}w)E_i^+(z)E_i^-(z)e^{\alpha_i}z^{H_{i,0}+1}E_j^+(w)E_j^-(w)e^{\alpha_j}w^{H_{j,0}+1} \\ &= \frac{z\kappa^{m_{ij}} - q^{-1}w}{z\kappa^{\frac{1}{2}m_{ij}}(1 - q^{-1}\kappa^{-m_{ij}}\frac{w}{z})}E_i^+(z)E_j^+(w)E_i^-(z)E_j^-(w) \\ &\quad \times e^{\alpha_i}e^{\alpha_j}z^{H_{i,0}+1}w^{H_{j,0}+1} \\ &= \kappa^{\frac{1}{2}m_{ij}}E_i^+(z)E_j^+(w)E_i^-(z)E_j^-(w)e^{\alpha_i}e^{\alpha_j}z^{H_{i,0}+1}w^{H_{j,0}+1}. \end{aligned}$$

On the other hand we have

$$\begin{aligned}
& (q^{\langle h_i, \alpha_j \rangle} \kappa^{m_{ij}} z - w) E_j(w) E_i(z) \\
&= \frac{q^{-1} \kappa^{m_{ij}} z - w}{w \kappa^{-\frac{1}{2} m_{ij}} (1 - q^{-1} \kappa^{m_{ij}} \frac{z}{w})} E_j^+(w) E_i^+(z) E_j^-(w) E_i^-(z) \\
&\quad \times e^{\alpha_j} e^{\alpha_i} w^{H_{j,0}+1} z^{H_{i,0}+1} \\
&= \kappa^{\frac{1}{2} m_{ij}} E_i^+(z) E_j^+(w) E_i^-(z) E_j^-(w) e^{\alpha_i} e^{\alpha_j} z^{H_{i,0}+1} w^{H_{j,0}+1}.
\end{aligned}$$

Thus we have $(z \kappa^{m_{ij}} - q^{\langle h_i, \alpha_j \rangle} w) E_i(z) E_j(w) = (q^{\langle h_i, \alpha_j \rangle} \kappa^{m_{ij}} z - w) E_j(w) E_i(z)$.

The formula $(z \kappa^{m_{ij}} - q^{-\langle h_i, \alpha_j \rangle} w) F_i(z) F_j(w) = (q^{-\langle h_i, \alpha_j \rangle} \kappa^{m_{ij}} z - w) F_j(w) F_i(z)$ is proved similarly.

Let us prove (A.1.12). Assume that $\langle h_i, \alpha_j \rangle = -1$. This is the most complicated case. The other cases can be proved similarly.

We have the following formulas:

$$\begin{aligned}
& E_i(z_1) E_i(z_2) E_j(w) \\
&= \frac{(z_1 - z_2)(z_1 - q^{-2} z_2) \kappa^{-m_{ij}}}{z_1 z_2 (1 - \frac{q^{-1} \kappa^{-m_{ij}} w}{z_1}) (1 - \frac{q^{-1} \kappa^{-m_{ij}} w}{z_2})} E_i^+(z_1) E_i^+(z_2) E_j^+(w) E_i^-(z_1) E_i^-(z_2) E_j^-(w) \\
&\quad \times e^{2\alpha_i} e^{\alpha_j} z_1^{H_{i,0}+1} z_2^{H_{i,0}+1} w^{H_{j,0}+1},
\end{aligned}$$

$$\begin{aligned}
& E_i(z_1) E_j(w) E_i(z_2) \\
&= \frac{(z_1 - z_2)(z_1 - q^{-2} z_2)}{z_1 w (1 - \frac{q^{-1} \kappa^{-m_{ij}} w}{z_1}) (1 - \frac{q^{-1} \kappa^{-m_{ij}} z_2}{w})} E_i^+(z_1) E_i^+(z_2) E_j^+(w) E_i^-(z_1) E_i^-(z_2) E_j^-(w) \\
&\quad \times e^{2\alpha_i} e^{\alpha_j} z_1^{H_{i,0}+1} z_2^{H_{i,0}+1} w^{H_{j,0}+1},
\end{aligned}$$

$$\begin{aligned}
& E_j(w) E_i(z_1) E_i(z_2) \\
&= \frac{(z_1 - z_2)(z_1 - q^{-2} z_2) \kappa^{m_{ij}}}{w^2 (1 - \frac{q^{-1} \kappa^{m_{ij}} z_1}{w}) (1 - \frac{q^{-1} \kappa^{m_{ij}} z_2}{w})} E_i^+(z_1) E_i^+(z_2) E_j^+(w) E_i^-(z_1) E_i^-(z_2) E_j^-(w) \\
&\quad \times e^{2\alpha_i} e^{\alpha_j} z_1^{H_{i,0}+1} z_2^{H_{i,0}+1} w^{H_{j,0}+1}.
\end{aligned}$$

Therefore it is enough to show that

$$\begin{aligned}
& \sum_{\sigma \in \mathfrak{S}_2} (z_{\sigma(1)} - z_{\sigma(2)})(z_{\sigma(1)} - q^{-2}z_{\sigma(2)}) \\
& \times \left\{ \frac{\kappa^{-m_{ij}}}{z_{\sigma(1)}z_{\sigma(2)}\left(1 - \frac{q^{-1}\kappa^{-m_{ij}}w}{z_{\sigma(1)}}\right)\left(1 - \frac{q^{-1}\kappa^{-m_{ij}}w}{z_{\sigma(2)}}\right)} - \frac{q + q^{-1}}{z_{\sigma(1)}w\left(1 - \frac{q^{-1}\kappa^{-m_{ij}}w}{z_{\sigma(1)}}\right)\left(1 - \frac{q^{-1}\kappa^{m_{ij}}z_{\sigma(2)}}{w}\right)} \right. \\
& \left. + \frac{\kappa^{m_{ij}}}{w^2\left(1 - \frac{q^{-1}\kappa^{m_{ij}}z_{\sigma(1)}}{w}\right)\left(1 - \frac{q^{-1}\kappa^{m_{ij}}z_{\sigma(2)}}{w}\right)} \right\} \\
& = 0.
\end{aligned} \tag{A.1.14}$$

This identity is proved by a direct calculation.

Thus the proposition is proved.

Acknowledgment . The author would like to thank Masaki Kashiwara, Tetsuji Miwa and Michio Jimbo for their encouragement and valuable discussions. He also thanks Kenji Iohara, Norio Suzuki, Eric Vasserot and Hiroshi Yamada for stimulating discussions.

REFERENCES

- [Be] J. Beck, *Braid group action and quantum affine algebras*, Commun. Math. Phys. 165 (1994), 555-568.
- [B] S. Berman, *On generators and relations for certain involutory subalgebras Kac-Moody Lie algebras*, Commun. Alg. 17(12) (1989), 3165-3185.
- [BM] S. Berman and R. Moody, *Lie algebras graded by finite root systems and the intersection matrix algebras of Slodowy*, Inv. Math. (1992), 323-347.
- [C] I. Cherednik, *Double-affine Hecke algebras, Knizhnik-Zamolodchikov equations and Macdonald's operators*, Intern. Math. Research Notices 9 (1992), 171-180.
- [CP] V. Chari and A. Pressly, *Quantum affine algebras and affine Hecke algebras*, Pacific Jour. of Math. 174 No.2 (1996), 295-326.
- [D1] V. Drinfeld, *Quantum groups*, Proc. ICM Berkeley 1 (1986), 789-820.
- [D2] V. Drinfeld, *A new realization of Yangians and quantized affine algebras*, Sov. Math. Dokl. 36 (1988), 212-216.
- [EM] S. Eswara Rao and R. V. Moody, *Vertex representations for N-toroidal Lie algebras and a Generalization of the Virasoro algebra*, Commun. Math. Phys. 159 (1994), 239-264.
- [FJ] I. Frenkel and N. Jing, *Vertex representations of quantum affine algebras*, Proc. Natl. Acad. Sci. USA 85 (1988), 9373-9377.
- [FK] I. Frenkel and V. Kac, *Basic Representations of affine Lie algebras and dual Resonance models*, Inv. Math. 62 (1980), 23-66.
- [GKV] V. Ginzburg, M. Kapranov and E. Vasserot, *Langlands reciprocity for algebraic surface*, Math. Res. Lrtt. 2 (1995), 147-160.
- [K] Y. Koyama, *Staggered polarization of vertex model with $U_q(\widehat{\mathfrak{sl}}(n))$ -symmetry*, Commun. Math. Phys. 164 (1994), 277-291.
- [MEY] R. V. Moody, S. Eswara Rao and T. Yokomuma, *Toroidal Lie algebras and vertex representations*, Geom. Dedicata 35 (1990), 283-307.

- [S] P. Slodowy, *Beyond Kac-Moody algebras, and inside*, Canadian Math. Soc. Conference Proceedings Vol.5 (1986), 361-371.
- [V] E. Vasserot, *Private communication*.
- [VV] M. Varagnolo and E. Vasserot, *Schur duality in the toroidal setting* Commun. Math. Phys. 182 (1996), 469-484.
- [Y] H. Yamada, *Extended affine Lie algebra and their vertex representations*, Publ. RIMS. 25 (1989), 587-603.

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY,
JAPAN